

## Know what to expect each week

Follow your progress with this chart. Please note, since everyone’s skin is different, results process at a different pace for different skin types. If you want to get the full effect of ReFissa, use it for at least 24 weeks.

WEEK			
<p>You may see redness, flaking and irritation of your skin. How do you know if this is a reaction to tretinoin (the active ingredient in ReFissa) or a sign it’s working? A good way to tell is by your own comfort level.</p> <p>It’s normal to experience some irritation during the first few months of treatment. If your skin becomes too irritated, consult with your doctor, who may suggest you try using ReFissa every other night.</p> <p>There are rare instances in which people with sensitive skin cannot tolerate the medication.</p>	<p>You may notice that your skin is smoother and has a softer texture.<sup>1</sup></p>	<p>Irregular pigmentation may begin to fade, while fine facial wrinkles may begin to diminish.</p> <p>You may see an overall improvement in the texture and appearance of your skin.<sup>1</sup></p>	<p>You may continue to see and feel your skin texture and skin tone improve.<sup>1</sup></p> <p>To maintain the results of ReFissa, continue to use it nightly.</p>

1. Renova® (tretinoin emollient cream) 0.05% product monograph. Ortho-McNeil Pharmaceuticals, Inc.; Printed 1996. ReFissa 0.05% is a prescription medicine that may reduce fine facial wrinkles and mottled hyperpigmentation in patients who also protect their skin from the sun and wear sunscreen daily. ReFissa does not eliminate wrinkles, repair sun damaged skin or reverse photo-aging. Do not use if you are pregnant, attempting pregnancy, or nursing. Do not use if taking medicines that may increase your sensitivity to sunlight. Use of ReFissa may make your skin more likely to burn from sunlight. ReFissa, early in treatment, may cause redness, itching, burning, stinging and peeling. If you have questions about side effects, contact your physician. If you are uncomfortable, use less medication and decrease the frequency of application. If discomfort is still significant, discontinue use and consult your physician. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088. Please see full Prescribing Information and Patient Package Insert. ReFissa is a trademark of Suneva Medical, Inc. © 2013 Suneva Medical, Inc. All rights reserved.